

A Wacky Graph of a Simple
Looking Function: The Problem

K. Ming Leung

Abstract: Plotting of a seemingly simple function can sometimes give unexpected wacky results that are due to the finiteness of the underlying floating-point system.

Directory

- [Table of Contents](#)
- [Begin Article](#)

Suppose we are given a function

$$f(x) = \frac{1 - \cos(x)}{x^2},$$

and we want to know how it looks like by plotting it for specific ranges of values for x . One very easy way is to use the built-in function called `ezplot` in MATLAB. This can be done by typing the following at the command prompt

```
ezplot('(1-cos(x))./(x.*x)', [0,10]);
```

The function was specified by the first argument of the `ezplot` function, and the second argument gives the range of x values for the plot. The plot looks great and it was certainly very easy to make indeed.

Suppose we are more interested in the range where x is small. We can shorten the range and plot again by entering the command

```
ezplot('(1-cos(x))./(x.*x)', [0,1e-2]);
```

Here the number `1e-2` means 1×10^{-2} . The resulting graph looks rather uninteresting. However on closer inspection we notice a small blip in the curve near $x = 0$. Actually MATLAB's `ezplot` function does not plot the end points of the given interval. It seems that the




Back



Doc






Back

◀ Doc


Doc ▶



blip cannot be due to the potential singularity of the function at $x = 0$. One can do a closer inspection of the blip by using the zoom feature in MATLAB's graphics window.

Re-plotting the function for an even small range of x values gives even wackier results. We want to find out what is really going on.


[Back](#)[◀ Doc](#)[Doc ▶](#)



Back

◀ Doc

Doc ▶



Back

◀ Doc

Doc ▶

First let us use calculus to find out the proper behavior of the function for x at the origin. When $x = 0$ both the numerator and denominator of the function go to zero, so we need to use the l'Hôpital's rule to get

$$\lim_{x \rightarrow 0} f(x) = \frac{\frac{d}{dx}(1 - \cos x)|_{x=0}}{\frac{d}{dx}(x^2)|_{x=0}} = \frac{\sin x|_{x=0}}{2x|_{x=0}}.$$

The numerator and denominator still both go to zero at $x = 0$, so we have use the l'Hôpital's rule again to get

$$\lim_{x \rightarrow 0} f(x) = \frac{\frac{d}{dx} \sin x|_{x=0}}{\frac{d}{dx} 2x|_{x=0}} = \frac{\cos x|_{x=0}}{2|_{x=0}} = \frac{1}{2}.$$


Therefore we see that the function must approach the constant value of $1/2$ at $x = 0$. Thus the results we are seeing here are highly incorrect.

To try to understand the cause of the problem we need to write a program to compute the function and analyze the results ourselves. We concentrate on the interval $[0, 1 \times 10^{-7}]$ and obtain the result as shown in the following graph. The MATLAB program, `wackyPlot0.m`



Back





is available on the course website. This result is clearly also incorrect.

The graph has a lot of prominent features and looks rather complicated. When we understand how real numbers are represented in a digital computer it turns out that we will be able to understand quantitatively every single feature exhibited in this graph.



Back

◀ Doc

Doc ▶